

bile collected for 24 h was treated with .beta.-glucuronidase and arylsulfatase at 37.degree. for 3 h to give I (R1 = Me, R2 = CO2Me) (III) and I (R1 = CO2Me, R2 = Me) (IV). III or IV was incubated with HL-60 human leukemia cells to increase the no. of cells capable of reducing NBT. Tablets and injections contg. III or IV were also formulated.


IT 221176-86-9P

RL: BAC (Biological activity or effector, except adverse); BMF (Bioindustrial manufacture); BPN (Biosynthetic preparation); MFM (Metabolic formation); THU (Therapeutic use); BIOL (Biological study); FORM (Formation, nonpreparative); PREP (Preparation); USES (Uses) (prodn. of active vitamin D derivs. from bile of animal given dihydroxyvitamin D4 and their uses)

RN 221176-86-9 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19)-trien-26-oic acid, 1,3,24,25-tetrahydroxy-, methyl ester, (1.alpha.,3.beta.,5Z,7E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



L11 ANSWER 6 OF 37 HCAPLUS COPYRIGHT 2002 ACS

AN 1999:106958 HCAPLUS

DN 130:209851

TI Preparation of active vitamin D derivatives and their use as bone density improvers, differentiation inducers, and immunosuppressants causing no hypercalcemia

IN Tachibana, Yoji

PA Nisshin Flour Milling Co., Ltd., Japan

SO Jpn. Kokai Tokkyo Koho, 10 pp.


CODEN: JKXXAF

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 11035553	A2	19990209	JP 1997-194200	19970718
OS	MARPAT 130:209851				
GI					



AB The derivs. I (R1, R2, R5 = H, OH; R1 = R2 .noteq. OH; R3, R4 = H, OH, Me; R3 = R4 .noteq. H, Me) are prepd. by irradiation of diene compds. II (R1-R5 = same as above) with UV ray, then thermal isomerization. II (R1 = R2 = R3 = H, R4 = Me, R5 = OH) (200 mg) was irradiated with a high-pressure Hg lamp for 10 min and subjected to thermal isomerization to give 24 mg I (R1 = R2 = R3 = H, R4 = Me, R5 = OH). The product showed high affinity to 1,25-dihydroxyvitamin D3 receptor.

IT 179189-36-7P 220875-68-3P


RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)


(prepn. of active vitamin D derivs. as bone d. improvers, differentiation inducers, and immunosuppressants causing no hypercalcemia)

RN 179189-36-7 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19)-triene-1,3,24,25-tetrol, (1.alpha.,3.beta.,5Z,7E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.





RE.CNT 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 13 OF 95 HCAPLUS COPYRIGHT 2002 ACS

AN 1999:502218 HCAPLUS

DN 131:284266

TI Physiological significance of C-28 hydroxylation in the metabolism of 1.alpha.,25-dihydroxyvitamin D2

AU Rao, D. Sunita; Siu-Caldera, Mei-Ling; Uskokovic, Milan R.; Horst, Ronald L.; Reddy, G. Satyanarayana

CS Department of Pediatrics, Women and Infants' Hospital of Rhode Island, Brown University School of Medicine, Providence, RI, 02905, USA

SO Arch. Biochem. Biophys. (1999), 368(2), 319-328

CODEN: ABBIA4; ISSN: 0003-9861

PB Academic Press

DT Journal

LA English

AB In our previous study, we indicated for the first time that C-28 hydroxylation plays a significant role in the metab. of 1.alpha.,25-dihydroxyvitamin D2 [1.alpha.,25(OH)2D2] by identifying 1.alpha.,24(S),25,28-tetrahydroxyvitamin D2 [1.alpha.,24(S),25,28(OH)4D2] as a major renal metabolite of 1.alpha.,25(OH)2D2 [G. S. Reddy and K-Y. Tserng, 1986]. The present study was performed to establish the physiol. significance of C-28 hydroxylation in the metab. of 1.alpha.,25(OH)2D2. We perfused rat kidneys in vitro with 1.alpha.,25(OH)2[26,27-3H]D2 (5 .times. 10-10M) and demonstrated that 1.alpha.,24(R),25-trihydroxyvitamin D2 [1.alpha.,24(R),25(OH)3D2] and 1.alpha.,24(S),25,28(OH)4D2 are the only two major physiol. metabolites of 1.alpha.,25(OH)2D2. In the same perfusion expts., we also noted that there is no conversion of 1.alpha.,25(OH)2D2 into 1.alpha.,25,28-trihydroxyvitamin D2[1.alpha.,25,28(OH)3D2]. Moreover, 1.alpha.,24(S),25,28(OH)4D2 is not formed in the perfused rat kidney when synthetic 1.alpha.,25,28(OH)3D2 is used as the starting substrate. This finding indicates that C-28 hydroxylation of 1.alpha.,25(OH)2D2 occurs only after 1.alpha.,25(OH)2D2 is hydroxylated at C-24 position. At present the enzyme responsible for the C-28 hydroxylation of 1.alpha.,24(R),25(OH)3D2 in rat kidney is not known. Recently, it was found that 1.alpha.,25(OH)2D3-24-hydroxylase (CYP24) can hydroxylate carbons 23, 24, and 26 of various vitamin D3 compds. Thus, it may be speculated that CYP24 may also be responsible for

the C-28 hydroxylation of 1.alpha.,24(R),25(OH)3D2 to form 1.alpha.,24(S),25,28(OH)4D2. The biol. activity of 1.alpha.,24(S),25,28(OH)4D2, detd. by its ability to induce intestinal calcium transport and bone calcium resorption in the rat, was found to be almost negligible. Also, 1.alpha.,24(S),25,28(OH)4D2 exhibited very low binding affinity toward bovine thymus vitamin D receptor. These studies firmly establish that C-28 hydroxylation is an important enzymic reaction involved in the inactivation of 1.alpha.,25(OH)2D2 in kidney under physiol. conditions. (c) 1999 Academic Press.


IT 100496-04-6

RL: BAC (Biological activity or effector, except adverse); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
(dihydroxyvitamin D2 C-28 hydroxylation and biol. inactivation by rat kidney)

RN 100496-04-6 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,24,25-tetrol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



RE.CNT 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 14 OF 95 HCAPLUS COPYRIGHT 2002 ACS

AN 1999:396342 HCAPLUS

DN 131:194395

TI Vitamin D assays and their clinical utility

AU Horst, Ronald L.; Hollis, Bruce W.

CS Metabolic Diseases and Immunology Research Unit, Agricultural Research Service, National Animal Disease Center, US Department of Agriculture, Ames, IA, USA

SO Vitam. D (1999), 239-271. Editor(s): Holick, Michael F.


Publisher: Humana, Totowa, N. J.

CODEN: 67UOAT

DT Conference; General Review

LA English

AB A review, with 139 refs., on assay methodol. for vitamin D and its metabolites 25-hydroxyvitamin D3, 24,25-dihydroxyvitamin D3, 1.alpha.,25-dihydroxyvitamin D3, 1,24,25-trihydroxyvitamin D3,



AB The title compd. I was prepd. from C22-aldehyde II (TBS = tert-butyldimethylsilyl) in 51% overall yield. The key feature of this synthesis is a one-pot construction of the requisite side chain using .alpha.-lithiomethylenetriphenylphosphorane $\text{Ph}_3\text{P:CHLi}$ and com. available (S)-2-methylglycidol III (TDS = thexyldimethylsilyl).


IT 103335-39-3P

RL: SPN (Synthetic preparation); PREP (Preparation)
(prepn. of)

RN 103335-39-3 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19),22-tetraene-1,3,25,26-tetrol,
(1.alpha.,3.beta.,5Z,7E,22E,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



L13 ANSWER 31 OF 95 HCAPLUS COPYRIGHT 2002 ACS
AN 1994:182476 HCAPLUS
DN 120:182476

TI metabolism and biological activity of 1,25(OH)2D2 and its metabolites in a chronic myelogenous leukemia cell line, RWLeu-4

AU Clark, J.W.; Reddy, G.S.; Santos-Moore, A.; Wankadiya, K.F.; Reddy, G.P.; Eil, C.; Lasky, S.; Tserng, K Y.; Horst, R.L.

CS Sch. Med., Brown Univ., Providence, RI, 02908, USA

SO Bioorg. Med. Chem. Lett. (1993), 3(9), 1873-8
CODEN: BMCLE8; ISSN: 0960-894X

DT Journal

LA English


AB The authors previously described the metab. of 1,25(OH)2D2 into various side chain hydroxylated metabolites formed as a result of C-24, C-26 and C-28 hydroxylations in rat kidneys. The authors now demonstrate C-24 hydroxylation of 1,25(OH)2D2 in human leukemic cells and also present evidence to show that C-24 and C-26 hydroxylations either alone or in combination do not significantly alter the effect of 1,25(OH)2D2 on these cells, while C-28 hydroxylation reduces its activity.

IT 100496-04-6 103305-11-9
RL: BPR (Biological process); BIOL (Biological study); PROC (Process) (metab. and antitumor activity of, in myelogenous leukemia of humans, as dihydroxyvitamin D2 metabolite)

RN 100496-04-6 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,24,25-tetrol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.




RN 103305-11-9 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,24,25,26-pentol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

L13, ANSWER 54 OF 95 HCAPLUS COPYRIGHT 2002 ACS
AN 1987:452924 HCAPLUS
DN 107:52924
TI Metabolic pathways from 1.alpha.,25-dihydroxyvitamin D3 to
1.alpha.,25-dihydroxyvitamin D3-26,23-lactone. Stereo-retained and
stereo-selective lactonization
AU Ishizuka, Seiichi; Norman, Anthony W.
CS Dep. Biochem., Teijin Inst. Bio-Med. Res., Hino, 191, Japan
SO J. Biol. Chem. (1987), 262(15), 7165-70
CODEN: JBCHA3; ISSN: 0021-9258
DT Journal
LA English
GI




AB Naturally occurring 23(S),25(R)-1.alpha.,25-dihydroxyvitamin D3 26,23-lactone (I) was produced in increasing amts. from 1.alpha.,25-dihydroxyvitamin D3 (II); 1.alpha.,25(R),26-trihydroxyvitamin D3 [1.alpha.,25(R),26-(OH)3D3]; 1.alpha.,25(S),25-(OH)3D3; 1.alpha.,23(S),25(R),26-tetrahydroxyvitamin D3 [1.alpha.,23(S),25(R),26-(OH)4D3]; and 23(S),25(R)-1.alpha.,25-dihydroxyvitamin D3 26,23-lactol (III) by II-supplemented chicken kidney and intestine mucosa homogenates. Thus, there are 2 possible metabolic pathways from II to 1.alpha.,23(S),25(R),26-(OH)4D3: the major pathway is by way of 1.alpha.,23(S),25-(OH)3D3 and the minor pathway is by way of 1.alpha.,25(R),26-(OH)3D3. 1.alpha.,23(S),25(R),26-(OH)4D3 is further metabolized to I via III. III was isolated in pure form and identified by UV spectrophotometry and mass spectrometry. Lactonization of 1.alpha.,23(S),25(R),26-(OH)4D3 and III occurred in a stereo-retained and stereoselective fashion.

IT 108131-93-7 108131-94-8 108211-12-7
RL: BIOL (Biological study)
(dihydroxyvitamin D3 lactone) formation from, by kidney, stereochem. in relation to)

RN 108131-93-7 HCAPLUS

CN 9,10-Secosterolesta-5,7,10(19)-triene-1,3,23,25,26-pentol,
(1.alpha.,3.beta.,5Z,7E,23R,25R)-(9CI) (CA INDEX NAME)

Double bond geometry as shown.




RN 108131-94-8 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,23,25,26-pentol,
(1.alpha.,3.beta.,5Z,7E,23R,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




RN 108211-12-7 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,23,25,26-pentol,
(1.alpha.,3.beta.,5Z,7E,23S,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.



IT 81176-40-1 100634-18-2

RL: BIOL (Biological study)


(dihydroxyvitamin D3 lactone formation from, in intestine and kidney)

RN 81176-40-1 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,25,26-tetrol,
(1.alpha.,3.beta.,5Z,7E,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




RN 100634-18-2 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,23,25,26-pentol,
(1.alpha.,3.beta.,5Z,7E,23S,25R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.



L13 ANSWER 58 OF 95 HCAPLUS COPYRIGHT 2002 ACS

AN 1986:587053 HCAPLUS

DN 105:187053

TI Isolation and identification of vitamin D metabolites

AU Napoli, Joseph L.; Koszewski, Nick J.; Horst, Ronald L.

CS Dep. Biochem., Univ. Buffalo, Buffalo, NY, 14214, USA

SO Methods Enzymol. (1986), 123(Vitam. Coenzymes, Pt. H), 127-40

CODEN: MENZAU; ISSN: 0076-6879

DT Journal

LA English

AB Extn. and cleanup of vitamin D metabolites from animal tissues and cells are briefly discussed. HPLC systems for sepg. hydroxylated vitamin D derivs. are examd. in detail; elution patterns of metabolites with various stationary and mobile phases are outlined. UV spectroscopy, electron-impact and chem.-ionization mass spectroscopy, chem. modification prior to detn., and NMR spectroscopy of vitamin D metabolites are presented.

IT 78780-98-0 100496-04-6 101151-55-7

RL: ANST (Analytical study)


(sepn. and detection of, by HPLC and spectroscopy)

RN 78780-98-0 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,25,26-tetrol,
(1.alpha.,3.beta.,5Z,7E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.


Double bond geometry as shown.



RN 100496-04-6 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,24,25-tetrol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



RN 101151-55-7 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,25,26-tetrol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.

achieved with analogs of I in accordance with their binding affinity for the hormone's receptor. Only cells with I receptor protein were inhibited in their colony formation by vitamin D analogs, indicating that the hormone receptor complex may be integrally involved in the in vitro suppression of the anchorage-independent phenotype.

IT 78780-98-0

RL: BIOL (Biological study)


(cancer cell growth inhibition by, receptor in relation to)

RN 78780-98-0 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,25,26-tetrol,
(1.alpha.,3.beta.,5Z,7E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.



L13 ANSWER 61 OF 95 HCAPLUS COPYRIGHT 2002 ACS

AN 1986:531100 HCAPLUS

DN 105:131100

TI Isolation and identification of 1,24,25-trihydroxyvitamin D2, 1,24,25,28-tetrahydroxyvitamin D2, and 1,24,25,26-tetrahydroxyvitamin D2: new metabolites of 1,25-dihydroxyvitamin D2 produced in the rat kidney

AU Reddy, G. Satyanarayana; Tserng, Kou Yi

CS Cleveland Metrop. Gen. Hosp., Case West. Reserve Univ., Cleveland, OH, 44109, USA


SO Biochemistry (1986), 25(18), 5328-36

CODEN: BICHAW; ISSN: 0006-2960

DT Journal

LA English

GI

I, R=R¹=HII, R=H, R¹=OHIII, R=OH, R¹=H

AB Three new metabolites of vitamin D₂ were produced in vitro by perfusing isolated rat kidneys with 1,25-dihydroxyvitamin D₂. They were isolated and purified from the kidney perfusate by the techniques of MeOH-CH₂Cl₂ lipid extn. and HPLC. By means of UV absorption spectrophotometry, mass spectrometry, and specific chem. reactions, the metabolites were identified as 1,24,25-trihydroxyvitamin D₂ (I) 1,24,25,28-tetrahydroxyvitamin D₂ (II), and 1,24,25,26-tetrahydroxyvitamin D₂ (III). Both II and III were also produced when a kidney was perfused with I. Thus, it becomes clear that 1,25-dihydroxyvitamin D₂ is 1st hydroxylated at C-24 to form I, which is then further hydroxylated at C-28 and C-26 to form II and III, resp. From several recent studies, it has been well established that 1,25-dihydroxyvitamin D₃ is converted into various further metabolites in the kidney as a result of chem. reactions such as C-23, C-24, and C-26 hydroxylations, C-24 ketonization, and C-23:C-26 lactonization. From this study it is obvious that 1,25-dihydroxyvitamin D₂ does not undergo all of the aforementioned chem. reactions except C-24 and C-26 hydroxylations. In addn., C-28 hydroxylation plays a significant role in the further metab. of 1,25-dihydroxyvitamin D₂.

IT 100496-04-6 103305-11-9


RL: FORM (Formation, nonpreparative)
(formation of, by kidney)

RN 100496-04-6 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,24,25-tetrol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




RN 103305-11-9 HCAPLUS

CN 9,10-Secoergosta-5,7,10(19),22-tetraene-1,3,24,25,26-pentol,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

Double bond geometry as shown.




IT 103321-15-9P


RL: SPN (Synthetic preparation); PREP (Preparation)
(prepn. and mass spectroscopy of)

RN 103321-15-9 HCAPLUS

CN 27-Nor-9,10-secoergosta-5,7,10(19),22-tetraen-25-one, 1,3,24-trihydroxy-,
(1.alpha.,3.beta.,5Z,7E,22E)- (9CI) (CA INDEX NAME)




L13 ANSWER 65 OF 95 HCAPLUS COPYRIGHT 2002 ACS
 AN 1986:108310 HCAPLUS
 DN 104:108310
 TI Biological activity and characteristics of 1.alpha.,25-(OH)2D3-26,23-lactone
 AU Ishizuka, S.; Kiyoki, M.; Orimo, H.; Norman, A. W.
 CS Teijin Inst. Bio-Med. Res., Tokyo, Japan
 SO Proc. Workshop Vitam. D (1985), 6th(Vitam. D), 402-3
 CODEN: PWVDDU; ISSN: 0721-7110
 DT Journal
 LA English
 GI




I

AB The biol. activity and characteristics of 4 diastereoisomers of 1.alpha.,25-dihydroxyvitamin D3 26,23-lactone (I) [75519-08-3] are discussed. The metabolic pathway for formation of the natural stereoisomer of I, 23(S),25(R)-I [81203-50-1] from 1.alpha.,25-dihydroxyvitamin D3 [32222-06-3] by chick kidney homogenate was through 1.alpha.,23(S),25(R),26-tetrahydroxyvitamin D3 [100634-18-2].



L13 ANSWER 68 OF 95 HCAPLUS COPYRIGHT 2002 ACS
 AN 1985:615623 HCAPLUS
 DN 103:215623
 TI (25S)-26-Hydroxycalcitriol
 CS Johann Wolfgang Goethe-Universitaet, Inst. Org. Chem., Frankfurt/Main,
 D-6000/50, Fed. Rep. Ger.
 SO Synform (1985), 3(2), 94-100
 CODEN: SNFMDF
 DT Journal; General Review
 LA English
 GI




I

AB Review of total syntheses of the title compd. (I) published since 1967.
 IT 77372-59-9P 78780-98-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (total synthesis of)
 RN 77372-59-9 HCAPLUS
 CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,25,26-tetrol,
 (1.alpha.,3.beta.,5Z,7E,25S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

Double bond geometry as shown.



RN 78780-98-0 HCAPLUS
 CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,25,26-tetrol,
 (1.alpha.,3.beta.,5Z,7E)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
 Double bond geometry as shown.



L13 ANSWER 69 OF 95 HCAPLUS COPYRIGHT 2002 ACS
 AN 1985:436524 HCAPLUS
 DN 103:36524
 TI Vitamin D metabolites regulate osteocalcin synthesis and proliferation of human bone cells in vitro
 AU Skjodt, H.; Gallagher, J. A.; Beresford, J. N.; Couch, M.; Poser, J. W.; Russell, R. G. G.
 CS Med. Sch., Univ. Sheffield, Sheffield, S10 2RX, UK
 SO J. Endocrinol. (1985), 105(3), 391-6
 CODEN: JOENAK; ISSN: 0022-0795
 DT Journal
 LA English
 AB The effects of 6 natural vitamin D metabolites of potential biol. and

therapeutic interest, 1,25-dihydroxyvitamin D3 (1,25-(OH)2D3) [32222-06-3], 25-hydroxyvitamin D3 (25-OH-D3) [19356-17-3], 24R,25-dihydroxyvitamin D3 (24R,25-(OH)2D3) [55721-11-4], 1,24R,25-trihydroxyvitamin D3 (1,24R,25-(OH)3D3) [56142-94-0], 25S,26-dihydroxyvitamin D3 (25S,26-(OH)2D3) [42737-59-7], and 1,25S,26-trihydroxyvitamin D3 (1,25S,26-(OH)2D3) [77372-59-9] on cell replication and expression of the osteoblastic phenotype in terms of osteocalcin prodn. were examd. in cultured human bone cells. At a dose of 5 .times. 10-12 mol/L, 1,25-(OH)2D3 stimulated cell proliferation, whereas at higher doses (5 .times. 10-9-5 .times. 10-6 mol/L) cell growth was inhibited in a dose-dependent manner. The same pattern of effects was seen for the other metabolites in a rank order of potency: 1,25-(OH)2D3 > 1,25S,26-(OH)3D3 = 1,24R,25-(OH)3D3 > 25S,26-(OH)2D3 = 24R,25-(OH)2D3 = 25-OH-D3. Synthesis of osteocalcin was induced by 1,25-(OH)2D3 in doses similar to those required to inhibit cell proliferation. Biphasic responses were obsd. for some of the metabolites in terms of osteocalcin synthesis, inhibitory effects becoming apparent at 5 .times. 10-6 mol/L. The cells did not secrete osteocalcin spontaneously. These results indicate that vitamin D metabolites may regulate growth and expression of differentiated functions of normal human osteoblasts.


IT 77372-59-9

RL: BIOL (Biological study)
(human bone cells proliferation and osteocalcin formation regulation by)

RN 77372-59-9 HCAPLUS

CN 9,10-Secocholesta-5,7,10(19)-triene-1,3,25,26-tetrol,
(1.alpha.,3.beta.,5Z,7E,25S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.
Double bond geometry as shown.



L13 ANSWER 70 OF 95 HCAPLUS COPYRIGHT 2002 ACS

AN 1985:132338 HCAPLUS

DN 102:132338

TI Stereoselective total synthesis of 1.alpha.,25S,26-trihydroxycholecalciferol

AU Wovkulich, P. M.; Barcelos, F.; Batcho, A. D.; Sereno, J. F.; Baggiolini, E. G.; Hennessy, B. M.; Uskokovic, M. R.


CS Chem. Res. Dep., Hoffmann-La Roche Inc., Nutley, NJ, 07110, USA

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

IT 77372-59-9P

RN 77372-59-9 HCAPLUS

Absolute stereochemistry.
Double bond geometry as shown.



DT Journal